

Soils Map - Tillable Acres





State: lowa
County: Scott
Location: 7-80N-3E
Township: Winfield
Acres: 72.54
Date: 7/9/2019







Soils data provided by USDA and NRCS.

Code	Soil	Acres	Percent	CSR2	Non-	Irr	*i	*i	*i	*i	*i Tall	CSR2**	CSR	*n	*n	*n	*n NCCPI	*n
	Description		of field	Legend	Irr Class *c	Class *c	Corn	Alfalfa	Soybeans	Bluegrass	Grasses			NCCPI	NCCPI Corn	NCCPI Small Grains	Soybeans	NCCF Cottor
41C	Sparta loamy fine sand, 5 to 9 percent slopes	20.88	28.8%		IVs	lle	124.8	3.5	36.2	2.2	3.7	34	25	50	50	36	28	
84	Clyde clay loam, 0 to 3 percent slopes	15.09	20.8%		llw		224	4.7	65	4	6.7	88	75	86	86	4	73	
41B	Sparta loamy fine sand, 2 to 5 percent slopes	10.31	14.2%		IVs	lle	140.8	3.9	40.8	2.5	4.2	39	40	50	50	37	28	
826	Rowley silt loam, 0 to 2 percent slopes	10.11	13.9%		lw		80	2.1	23.2	1.4	2.4	76	95	87	87	51	83	
175B	Dickinson fine sandy loam, 2 to 5 percent slopes	6.45	8.9%		Ille		80	2.2	23.2	1.4	2.4	50	55	60	60	47	36	
41D	Sparta loamy fine sand, 9 to 14 percent slopes	4.21	5.8%		VIs		105.6	3	30.6	1.9	3.2	8	15	48	48	35	26	
961	Ambraw clay loam, 0 to 2 percent slopes	3.98	5.5%		llw		196.8	4.1	57.1	3.5	5.9	73	47	91	91	13	76	



977	Richwood silt loam, 0 to 2 percent slopes	1.15	1.6%		I		240	6.7	69.6	4.3	7.2	100	95	92	92	73	86	
430	Ackmore silt loam, 0 to 2 percent slopes	0.36	0.5%		llw		203.2	5.3	58.9	3.7	6.1	77	83	87	87	56	76	
Weighted Average						142.5	3.6	41.3	2.5	4.3	55.1	52	*n 66.5	*n 66.5	*n 31.9	*n 49.4	*n 0	

^{**}IA has updated the CSR values for each county to CSR2.

*i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

*n: The aggregation method is "Weighted Average using major components"

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.