





NE DAVIS EAST, LLC - 186.41 ACRES M/L (186.40 AC TOTAL, 122.02 AC TILLABLE)

## CRP SOILS



County: **Davis County, IA**  
Location: **18-70N-12W**  
Township: **Salt Creek**  
Acres: **116.36**  
Date: **02/09/2026**



Field borders obtained from Farm Service Agency  
as of 2008

FarmWorth, LLC makes no representations or warranties, express or implied, as to the accuracy of any information, data, numerical values, boundaries, or any other information generated through the use of FarmWorth.com. User is solely responsible for independently investigating and determining all information provided through FarmWorth.com prior to use and waives any and all claims against FarmWorth, LLC for any inaccuracies or inconsistencies in the information and/or data.

| Code     | Description                                                          | Acres | % of field | IA CSR2 | Soil Class (Non-Irr) | Drainage Class          |
|----------|----------------------------------------------------------------------|-------|------------|---------|----------------------|-------------------------|
| 425D2    | Keswick loam, 9 to 14 percent slopes, moderately eroded              | 25.43 | 21.85 %    | 9       | 4                    | Somewhat poorly drained |
| 132C2    | Weller silty clay loam, 5 to 9 percent slopes, moderately eroded     | 22.25 | 19.12 %    | 59      | 3                    | Moderately well drained |
| 424E2    | Lindley-Keswick loams, 14 to 18 percent slopes, moderately eroded    | 18.20 | 15.64 %    | 24      | 6                    | Well drained            |
| 425D3    | Keswick clay loam, 9 to 14 percent slopes, severely eroded           | 11.29 | 9.70 %     | 10      | 6                    | Somewhat poorly drained |
| 730B     | Nodaway-Cantril complex, 0 to 5 percent slopes                       | 11.12 | 9.55 %     | 77      | 2                    | Moderately well drained |
| 425C2    | Keswick loam, 5 to 9 percent slopes, moderately eroded               | 9.08  | 7.80 %     | 27      | 3                    | Somewhat poorly drained |
| 132B     | Weller silt loam, 2 to 5 percent slopes                              | 8.86  | 7.61 %     | 67      | 3                    | Moderately well drained |
| 594D2    | Galland loam, heavy loess, 9 to 14 percent slopes, moderately eroded | 3.02  | 2.59 %     | 19      | 4                    | Somewhat poorly drained |
| 795D2    | Ashgrove silty clay loam, 9 to 14 percent slopes, moderately eroded  | 2.78  | 2.39 %     | 7       | 4                    | Poorly drained          |
| 65E2     | Lindley loam, 14 to 18 percent slopes, moderately eroded             | 2.47  | 2.12 %     | 29      | 6                    | Well drained            |
| 1715     | Nodaway-Lawson-Ackmore silt loams, 0 to 2 percent slopes             | 1.41  | 1.21 %     | 78      | 3                    | Somewhat poorly drained |
| 65E      | Lindley loam, 14 to 18 percent slopes                                | 0.47  | 0.40 %     | 33      | 6                    | Well drained            |
| Average: |                                                                      |       |            | 34.9    |                      |                         |

IA has updated the CSR values for each county to CSR2.

Soils data provided by USDA and NRCS.