





# TRACT 10 TILLABLE SOILS



County: **Fayette County, IA**  
Location: **26-93N-9W**  
Township: **Center**  
Acres: **126.87**  
Date: **09/14/2026**



FarmWorth, LLC makes no warranties, express or implied, regarding the accuracy of any information provided through FarmWorth.com. User is solely responsible for independently verifying all information prior to use and waives any claims against FarmWorth, LLC arising from inaccuracies in such information.

| Code     | Description                                                       | Acres | % of field | IA CSR2 | Soil Class (Non-Irr) | Drainage Class               |
|----------|-------------------------------------------------------------------|-------|------------|---------|----------------------|------------------------------|
| 171B     | Bassett loam, 2 to 5 percent slopes                               | 39.32 | 30.99 %    | 85      | 2                    | Moderately well drained      |
| 198B     | Floyd loam, 1 to 4 percent slopes                                 | 34.76 | 27.40 %    | 89      | 2                    | Somewhat poorly drained      |
| 213B     | Rockton loam, 30 to 40 inches to limestone, 2 to 5 percent slopes | 12.39 | 9.77 %     | 53      | 2                    | Well drained                 |
| 409B     | Dickinson fine sandy loam, till substratum, 2 to 5 percent slopes | 7.62  | 6.01 %     | 74      | 3                    | Well drained                 |
| 284B     | Flagler sandy loam, 2 to 5 percent slopes                         | 5.89  | 4.64 %     | 49      | 3                    | Somewhat excessively drained |
| 177      | Saupe loam, 0 to 2 percent slopes                                 | 5.38  | 4.24 %     | 60      | 2                    | Well drained                 |
| 302C     | Coggon loam, 5 to 9 percent slopes                                | 4.73  | 3.73 %     | 71      | 3                    | Moderately well drained      |
| 84       | Clyde clay loam, 0 to 3 percent slopes                            | 3.48  | 2.74 %     | 88      | 2                    | Poorly drained               |
| 175B     | Dickinson fine sandy loam, 2 to 5 percent slopes                  | 3.02  | 2.38 %     | 50      | 3                    | Well drained                 |
| 83B      | Kenyon loam, 2 to 5 percent slopes                                | 2.38  | 1.87 %     | 90      | 2                    | Moderately well drained      |
| 391B     | Clyde-Floyd complex, 1 to 4 percent slopes                        | 1.96  | 1.54 %     | 87      | 2                    | Poorly drained               |
| 407B     | Schley loam, 1 to 4 percent slopes                                | 1.94  | 1.53 %     | 81      | 2                    | Somewhat poorly drained      |
| 171C     | Bassett loam, 5 to 9 percent slopes                               | 1.67  | 1.31 %     | 80      | 3                    | Moderately well drained      |
| 221      | Klossner muck, 1 to 4 percent slopes                              | 1.55  | 1.22 %     | 48      | 3                    | Very poorly drained          |
| 178B     | Waukee loam, 2 to 5 percent slopes                                | 0.39  | 0.31 %     | 64      | 2                    | Well drained                 |
| 302B     | Coggon loam, 2 to 5 percent slopes                                | 0.28  | 0.22 %     | 76      | 2                    | Moderately well drained      |
| 115F     | Sandy escarpments, 14 to 30 percent slopes                        | 0.12  | 0.09 %     | 5       | 7                    | Well drained                 |
| Average: |                                                                   |       |            | 77.7    |                      |                              |

